

List of pathogens with field resistance towards QoL fungicides (updated 12/09/12)

Species name		Host	Geographical distribution	Type of resistance	Citation
<i>Alternaria alternata</i> , <i>Alternaria tenussima</i> , <i>Alternaria arborescens</i>	Alternaria Blight	Pistachio	USA	G143A	(1) (2)
<i>Alternaria alternata</i>	Early Blight	Potato and Tomato	EU	G143A	(17)
<i>Alternaria mali</i>	Alternaria Blotch	Apple	USA	G143A	(3)
<i>Alternaria solani</i>	Early Blight	Potato	USA	F129L	(4)
<i>Ascochyta rabiei</i>	Ascochyta blight	chickpeas	USA	G143A	(32)

<i>Blumeria graminis</i> f. sp. <i>tritici</i> and <i>hordei</i>	Powdery Mildew	Wheat and Barley	EU	G143A	(5)
<i>Botrytis cinerea</i>	Gray Mold	Strawberries	EU	G143A	(17)
<i>Cercospora sojina</i>	Frogeye spot	Soybeans	USA	G143A?	(29)
<i>Cercospora beticola</i>	Cercospora Leaf spot	Sugar beet	USA	G143A	(31)
<i>Cladosporium carpophilum</i>	Scab	Almonds	USA	?	(27)
<i>Colletotrichum graminicola</i>	Anthracnose	Turf grass	USA	G143A	(6)
<i>Corynespora cassiicola</i>	Leaf Spot, Target Spot	Cucumber	Japan	G143A	(7)
<i>Didymella bryoniae</i>	Gummy Stem Blight	Cucurbit	USA	G143A	(8)
<i>Erysiphe necator</i>	Powdery Mildew	Grapes	USA, EU	G143A	(17)
<i>Glomerella cingulata</i> (= <i>Colletotrichum gloeosporioides</i>)	Anthracnose	Strawberries	Japan	G143A	(9)

<i>Microdochium nivale</i> <i>Microdochium majus</i>	Snow mold, Ear blight	Wheat	EU, Japan	G143A	(17, 28)
<i>Mycosphaerella fijiensis</i>	Black Sigatoka	Banana	Central and South America, CAM, PHIL	G143A	(10)
<i>Mycosphaerella graminicola</i>	Septoria leaf spot	Wheat	EU	G143A	(11) (12)
<i>Mycosphaerella musicola</i>	Yellow Sigatoka	Banana	S America, Australia	G143A	(17)
<i>Mycovellosiella nattrassii</i>	Leaf Mold	Eggplant	Japan	G143A	(13)
<i>Passalora fulva</i>	Leaf Mold	Tomato	Japan	F129L	()
<i>Phaeosphaeria nodorum</i>	S. nodorum leaf blotch	Wheat	EU	G143A	(30)
<i>Plasmopara viticola</i>	Downy Mildew	Grape	EU	G143A and F129L	(14)
<i>Pseudoperonospora cubensis</i>	Downy Mildew	Cucurbits	EU, Asia	G143A	(15) (16)
<i>Pyrenophora teres</i>	Net Blotch	Barley	EU	F129L	(17, 24, 26)

<i>Pyrenophora. Tritici-repentis</i>	Tan Spot	Wheat	EU	G143A, and F129L and G137R	(17, 24, 25)
<i>Pyricularia grisea</i>	Gray Leaf Spot	Turf grass	USA	G143A and F129L	(18) (19)
<i>Pythium aphanidermatum</i>	Pythium Blight	Turf grass	USA	F129L	(20)
<i>Ramularia areola</i>	Grey mildew	Cotton	Brazil	?	(17)
<i>Ramularia collo-cygni</i>	Ramulariose	Barley	EU	G143A	(17)
<i>Rhynchosporium secalis</i>	Scald	Barley	EU	G143A	(17)
<i>Rhizoctonia solani AG1.1A</i>	Sheath blight	Rice	USA	F129L	(17)
<i>Sphaerotheca fuliginea</i>	Powdery Mildew	Cucurbits	EU, Asia	G143A	(21) (22)
<i>Stemphylium vesicarium</i>	Brown spot	Asparagus and Pear	EU	G143A	(17)
<i>Venturia inaequalis</i>	Scab	Apple	EU	G143A	(23)

<i>Venturia pirina</i>	Scab	Pears	USA	G143A	(17)
------------------------	------	-------	-----	-------	------

Citations

- (1) Ma, Z., Felts, D. and Michailides, T.J. (2003) Resistance to azoxystrobin in *Alternaria* isolates from pistachio in California. *Pesticide Biochemistry and Physiology* 77: 66-74
- (2) Ma, Z. and Michailides, T. J. (2004) An allele-specific PCR assay for detecting azoxystrobin-resistant *Alternaria* isolates from pistachio in California. *Journal of Phytopathology* 152: 118-121
- (3) Lu, Y.L., Sutton, T.B. and Ypema, H. (2003) Sensitivity of *Alternaria mali* from North Carolina apple orchards to pyraclostrobin and boscalid. *Phytopathology* 93: S54 (Abstract)
- (4) Pasche, J.S., Wharam, C.M. and Gudmestad, N.C. (2002) Shift in sensitivity of *Alternaria solani* (potato early blight) to strobilurin fungicides. *Proceedings of the BCPC Conference Pests & Diseases 2002*: 841-846
- (5) Sierotzki H, Wullschleger J. and Gisi U. (2000) Point-mutation in cytochrome b gene conferring resistance to strobilurin fungicides in *Erysiphe graminis* f. sp. *Tritici* field isolates. *Pesticide Biochemistry and Physiology* 68:107-112.
- (6) Avila-Adame, C., Olaya, G. and Köller, W. (2003) Characterization of *Colletotrichum graminicola* isolates resistant to strobilurin-related QoI fungicides. *Plant Disease* 87: 1426-1432
- (7) Ishii H. (2004) Fungicide resistance: a factor limiting integrated disease control. Proceedings of the 15th international plant protection congress (ed Guo Yu-yuan) p 216
- (8) Langston D. (2002) Quadris Resistance in Gummy Stem Blight Confirmed. *Georgia Extension Vegetable News* 2 (1): 1-2
- (9) Ishii H. (2004) Fungicide resistance: a factor limiting integrated disease control. Proceedings of the 15th international plant protection congress (ed Guo Yu-yuan) p 216
- (10) Sierotzki H., Parisi S., Steinfeld U., Tenzer I., Poirey S. and GisiU. (2000) Mode of resistance to respiration inhibitors at the cytochrome bc1 complex of *Mycosphaerella fijiensis*. *Pest Management Science* 56: 833-841.
- (11) Fraaije B. A., Brunett F. J., Clark W. S., Motteram J., Lucas J. A. (2005 submitted) . In: Modern fungicides and antifungal compounds II, eds Lyr H., Russell P. E., Dehne H-W. Gisi U. Kuck K-H, 14th International Reinhardtsbrunn Symposium, AgroConcept, Bonn, Verlag Th. Mann Gelsenkirchen, pp
- (12) Sierotzki H., Pavic L., Hugelshofer U., Stanger C., Cleere S., Windass J. and Gisi U. (2005 submitted) Population dynamics of *Mycosphaerella graminicola* in response to selection by different fungicides. . In: Modern fungicides and antifungal compounds II, eds Lyr H., Russell P. E., Dehne H-W. Gisi U. Kuck K-H, 14th International Reinhardtsbrunn Symposium, AgroConcept, Bonn, Verlag Th. Mann Gelsenkirchen, pp

- (13) Ishii H. (2004) Fungicide resistance: a factor limiting integrated disease control. Proceedings of the 15th international plant protection congress (ed Guo Yu-yuan) p 216
- (14) Heaney S. P., Hall A. A., Davis S. A. and Olaya G. (2000) Resistance to fungicides in the QoI-STAR cross resistance group: current perspectives. In Proceedings Brighton Crop Protection Conference- Pests and Diseases 2: 755-762..
- (15) Heaney S. P., Hall A. A., Davis S. A. and Olaya G. (2000) Resistance to fungicides in the QoI-STAR cross resistance group: current perspectives. In Proceedings Brighton Crop Protection Conference- Pests and Diseases 2: 755-762..
- (16) Ishii H., Fraaije B A., Sugiyama T., Noguchi K., Nishimura K., Takeda T., Amano T. and Hollomon D. W. (2001) Occurrence and molecular characterization of strobilurin resistance in cucumber powdery mildew and downy mildew. *Phytopathology* 91: 1166-1171.
- (17) FRAC
- (18) Vincelli, P. and Dixon E. (2002) Resistance to QoI (Strobilurin-like) fungicides in isolates of *Pyricularia grisea* from perennial ryegrass. *Plant Disease* 86: 235-240
- (19) Kim, Y.S., Dixon, P., Vincelli, P. and Farman, M.L. (2003) Field resistance to strobilurin (QoI) fungicides in *Pyricularia grisea* caused by mutations in the mitochondrial cytochrome b gene. *Phytopathology* 93: 891-900
- (20) Gisi U., Sierotzki H., Cook A., McCaffery A. (2002) Mechanisms influencing the evolution of resistance to Qo inhibitor fungicides. Pest Management Science 58(9): 859 - 867
- (21) Heaney S. P., Hall A. A., Davis S. A. and Olaya G. (2000) Resistance to fungicides in the QoI-STAR cross resistance group: current perspectives. In Proceedings Brighton Crop Protection Conference- Pests and Diseases 2: 755-762..
- (22) Ishii H., Fraaije B A., Sugiyama T., Noguchi K., Nishimura K., Takeda T., Amano T. and Hollomon D. W. (2001) Occurrence and molecular characterization of strobilurin resistance in cucumber powdery mildew and downy mildew. *Phytopathology* 91: 1166-1171.
- (23) Steinfeld U., Sierotzki H., Parisi S. and Gisi U. (2002) Comparison of resistance mechanisms to strobilurin fungicides in *Venturia inaequalis*. In: Modern fungicides and antifungal compounds II, eds Lyr H., Russell P. E., Dehne H-W. Gisi U. Kuck K-H, 13th International Reinhardtsbrunn Symposium, AgroConcept, Bonn, Verlag Th. Mann Gelsenkirchen, pp. 167-176
- (24) Sierotzki H., Frey R., Wullschleger J., Palermo S., Karli S., Godwin J. and Gisi U. (2006, accepted for publication) Cytochrome b gene sequence and structure of *Pyrenophora teres* and *P. tritici-repentis* and implications for QoI resistance. Pest Management Science 63:225–233 (2007)
- (25) Stammmer G., Strobel D., Semar M. and Klappach K. (2006): Diagnostics of fungicide resistance and relevance of laboratory data for the field. *Aspects of Applied Biology* 78: 29-36
- (26) Semar M., Strobel D., Koch A. Klappach K. and Stammmer G. (2007): Field efficacy of pyraclostrobin against populations of *Pyrenophora teres* containing the F129L mutation in the cytochrome b gene. *Journal of Plant Diseases and Protection*: 114: 117-119
- (27) Foerster H, Connell J H, Adaskaveg J E (2009). QoI resistance in *Fusicladium carpophilum* populations from almond in California. *Phytopathology* 99 S35
- (28) First occurrence of resistance to strobilurin fungicides in *Microdochium nivale* and *Microdochium majus* from French naturally infected wheat grains. Walker A S, Auclair C, Gredt M, Leroux P. 2009 Pest management Science 65 906-915
- (29) Frogeye Leaf Spot Pathogen with Reduced Sensitivity to Fungicides Found in Tennessee Soybean Field.
<http://bulletin.ipm.illinois.edu/article.php?id=1434>
- (30) Blixt E, Djurle A, Yuen J, Olson Å (2009). Fungicide sensitivity in Swedish isolates of *Phaeosphaeria nodorum*. *Plant Pathology* 58, 655-664
- (31) Bolton M D, Rivera V, Secor G (2012). Identification of the G143A mutation associated with QoI resistance in *Cercospora beticola* field isolates

from Michigan, United States. Pest Management Science doi: 10.1002/ps.3358

(32) Delgado J. A., Lynnes T. C., Meinhardt S. W., Wise K. A., Gudmestad N. C., Bradley C. A., Markell S. G. and Goswami R. S. (2012). Identification of the mutation responsible for resistance to QoI fungicides and its detection in *Ascochyta rabiei* (teleomorph *Didymella rabiei*). Plant Pathology (2012) Doi: 10.1111/j.1365-3059.2012.02661.x